



# Salipro technology in membrane protein research

Siavash Mostafavi<sup>1,2</sup>, Tânia F. Custódio<sup>1,2</sup>,  
Katharina E. J. Jungnickel<sup>1,2</sup> and Christian Löw<sup>1,2,3</sup>

Reconstitution and direct extraction of membrane proteins using saposins is an emerging technique for solubilizing and stabilizing membrane proteins. The Salipro technology offers several advantages over traditional detergent solubilization, including a more native lipid environment, increased protein stability, and maintenance of functionality. This review covers recent studies that have used Salipros to characterize membrane proteins, as well as advances in direct extraction methods that have enabled the structural and functional characterization of a variety of targets.

## Addresses

<sup>1</sup> Centre for Structural Systems Biology (CSSB), Notkestraße 85, 22607 Hamburg, Germany

<sup>2</sup> European Molecular Biology Laboratory (EMBL) Hamburg, Notkestraße 85, 22607 Hamburg, Germany

<sup>3</sup> Bernhard Nocht Institute for Tropical Medicine, 20359 Hamburg, Germany

Corresponding authors: Mostafavi, Siavash ([siavash.mostafavi@embl-hamburg.de](mailto:siavash.mostafavi@embl-hamburg.de)); Löw, Christian ([christian.loew@embl-hamburg.de](mailto:christian.loew@embl-hamburg.de))  
✉ (Löw C.)

Current Opinion in Structural Biology 2025, 93:103050

This review comes from a themed issue on **VSI: Membranes (2025)**

Edited by **Scott Prosser** and **Liguo Wang**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online xxx

<https://doi.org/10.1016/j.sbi.2025.103050>

0959-440X/© 2025 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

## Introduction


Integral membrane proteins (IMPs) are essential to many biological processes but are notoriously difficult to study due to their hydrophobic nature and the challenges in isolating them in a functional, water-soluble state. To overcome these challenges, various membrane protein reconstitution systems have been developed [1], with the Salipro technology emerging as a highly promising and versatile approach. The Salipro membrane mimetic system (also referred to as Saposin nanoparticles and Saposin nanodiscs) has been established by Frauenfeld and colleagues [2] and uses the

mammalian lipid-binding protein saposin A (SapA) to stabilize and reconstitute IMPs into a lipid bilayer-like environment.

Saposins belong to a family of small glycoproteins that play a critical role in lipid metabolism, particularly in the breakdown of sphingolipids, which are key components of cell membranes. They are synthesized as part of a larger precursor protein called prosaposin (Uniprot: P07602), that is later processed into the individual saposins (SapA, SapB, SapC, and SapD), each of which is associated with specific enzymes that facilitate the degradation of complex lipids in the lysosome. Earlier work has already demonstrated that saposins can exist in different conformations [3]. In the absence of lipids, SapA, a nine kilodalton fragment of prosaposin spanning residues 60-140, adopts a closed monomeric conformation, while the structure opens in the presence of detergents or lipids and exposes a hydrophobic surface (Figure 1a). The X-ray structure of such a complex revealed two chains of SapA encapsulating 40 bound detergent molecules organized in an ordered bilayer-like hydrophobic core [4]. This lipoprotein structure is mainly stabilized by lipid self-assembly. Later, such an assembly was further confirmed with branched detergents such as lauryl maltose neopentyl glycol (LMNG) [5].

The Salipro system is based on these binding properties of SapA. SapA serves as a flexible scaffold that surrounds an IMP embedded in a small lipid bilayer patch (Figure 1b). This flexibility eliminates the need to screen multiple scaffolds. The membrane mimetic structure allows IMPs to maintain their native conformation, a critical feature for structural biology, drug discovery, and functional assays. The classical workflow for reconstituting IMPs into Salipros is similar to that of other systems, such as nanodiscs, where a detergent-purified membrane protein is combined with SapA and phospholipids. After detergent removal, the resulting Salipro can be purified and analysed to confirm successful reconstitution (Figure 1). But recent advances in this technology now allow the reconstitution of IMPs into Salipros while keeping the endogenous lipids bound [6]. Here we review the latest developments of this technique, highlight applications and provide examples where this technology led to new insights into the structure and function of IMPs.

Figure 1



(a) Schematic diagram of SapA binding to lipids. Upon the addition of lipids, SapA undergoes a conformational change to reveal the hydrophobic core, which then binds to lipid molecules. (b) highlights the general process of Salipro reconstitution. IMPs are either first solubilized and purified in detergent, then mixed with SapA and lipids. After detergent removal, SapA and lipids combine to form a scaffold around the IMP. Alternatively, membranes are partially solubilized using Digitonin or mild detergents, followed by the addition of SapA. In this scenario SapA forms a small belt around the IMPs bound to its endogenous lipids.


### Applications of Salipros for structure determination by cryo electron microscopy

Among the advantages offered by Salipros are preservation of bound lipids [7–11], improved protein stability [12,13], stabilization of oligomeric [14,15] or conformational states [10,16] and improved structure resolution by cryo electron microscopy (cryo-EM) [17]. All properties that make this reconstitution technology attractive for structure determination. Within the last few years, there have been almost a hundred Salipro-reconstituted structures deposited in the Protein Data Bank (Figure 2, Supplementary Table S1). Of these, about 25 % are unique structures, while the rest represent structures with different ligands or conformational states. A small gallery of selected structures is presented in Figure 2, and a subset where the SapA density is resolved are shown in Figure 3, highlighting that this technology can be applied to IMPs of various sizes, oligomeric

states and origins. The saposin density in these structures shows that SapA molecules arrange in a tilted array around the IMP and lipid belt (Figure 3).

Several of these structures highlight well-resolved densities corresponding to lipid molecules tightly interacting with the IMP. These bound lipids can alter the activity or conformation of the protein, sometimes revealing previously unknown states or even mediating ligand binding. For example, the reconstitution of the ACA8 Calcium Pump from *Arabidopsis thaliana* enabled the structural characterization of lipid-mediated regulation [18]. A series of structures of the human ligand-gated ion channel  $\alpha 1\beta 2\gamma 2$  GABA<sub>A</sub> receptor reconstituted in Salipros and bound to various ligands show well-resolved lipid densities [7,8,19,20]. In some cases, the lipids appear to be involved in modulating neurosteroid binding [8], while in other cases, such as when the receptor is bound to methaqualone and its analog 2-

Figure 2




**(a)** The cumulative number of publications utilizing Salipros for structural studies, as well as the cumulative number of structures published every year which shows an accelerating growth in the use of saposin reconstitution. **(b)** The choice of lipids for SapA reconstituted structures within the last years. **(c)** A gallery of selected, recently published structures of IMPs in Salipros. The density corresponding to the scaffold protein SapA and lipids has been filtered to aid in visualization.

phenyl-3-(p-tolyl)quinazolin-4(3H)-one (PPTQ), the lipids stabilize the bound drug molecule [7].

In the case of the human  $\rho 1$  GABA<sub>A</sub> receptor, endogenous neurosteroids such as  $\beta$ -estradiol and pregnenolone sulfate can modulate receptor response to GABA binding [21]. A structure of this receptor reconstituted in Salipros and bound to  $\beta$ -estradiol revealed a previously unknown binding pocket for the neurosteroid at the interface between the extracellular domain (ECD) and

the transmembrane domain (TMD) [21]. Interestingly,  $\beta$ -estradiol binding did not induce any significant conformational changes, however the presence of  $\beta$ -estradiol blocked allosteric conformational changes that link GABA binding at the ECD and pore opening within the TMD [21]. This inhibitory mechanism appears to be specific to  $\rho$  subtypes of the GABA<sub>A</sub> receptor. In contrast, the mechanism of pregnenolone sulfate-induced inhibition was shown to occur via binding within the pore and inhibition of ion permeation, which

Figure 3



(a, b) Examples of structures where the SapA scaffold protein is well-resolved, highlighting the alpha-helical organization of the scaffold protein, as well as demonstrating the low degree of flexibility within the scaffold. (c) By low-pass filtering the density corresponding to the disc, we were able to find additional examples of structures where the SapA scaffold is well-resolved.

is similar to the inhibitory mechanism of other steroids in the  $\alpha 1\beta 2\gamma 2$  GABA<sub>A</sub> receptor [21].


Lipids are known to affect the equilibrium and transitions between the various conformational states of nicotinic acetylcholine receptors (nAChR) [22], and a structure of the nAChR from the electric ray *Torpedo torpedo* in Salipros revealed a new lipid binding site within a helical bundle previously hypothesized to be important for lipid sensing by the receptor [23]. Similarly, in a structure of the protein Smoothened, a component of the hedgehog signalling pathway, reconstituted in Salipros and representing an inactive state, a previously unknown sterol binding site could be identified, as well as accompanying conformational differences compared to previous structures that may have important consequences for the hedgehog signalling pathway [10].

In addition to the presence of specifically bound lipids, a lipidic environment can have additional effects on the overall conformation of a protein and may stabilize

physiologically more relevant states compared to detergent. An example of this is the *Escherichia coli* multidrug efflux pump AcrB. Binding of the small protein AcrZ is known to alter the transporter's substrate selectivity [24]. A structure of the complex determined in Salipros revealed the structural mechanism of how cardiolipin and AcrZ cooperate to allosterically modulate the transporter's selectivity [24]. Notably, the structure differs from previously determined detergent solubilized crystal structures of the complex, and showed a strong bending of AcrZ towards the binding groove of AcrB (Figure 4a).

In the case of the murine 5-HT<sub>3A</sub> serotonin receptor in Salipros, its apo form adopts a more tightly packed 'coupled' state that extends to the boundary between the extracellular and transmembrane domains, compared to the detergent-solubilized structure [25] (Figure 4b). This region contains highly conserved residues important for gating. In addition, due to the lipid bilayer-like environment, the receptor shows a widened

Figure 4



Comparisons of detergent and Salipro reconstituted structures. **(a)** In the AcrBZ complex, most of the AcrB subunit does not show large differences (upper panel), but the AcrZ subunit is more bent towards the membrane in the Salipro reconstituted structure. **(b)** There are major conformational differences within the transmembrane domains (lower panel), but not in the extracellular part of the 5-HT<sub>3A</sub> receptor (upper panel) when solubilized in detergent compared to the Salipro reconstituted receptor. **(c)** Reconstitution of the 5-HT<sub>3A</sub> receptor in Salipros also revealed an additional tetrameric, as opposed to pentameric, assembly state.

open pore upon serotonin binding compared to the detergent-solubilized structure [25]. The saposin-reconstituted receptor bound to serotonin is also present in an asymmetric form, speculated by the authors to be an intermediate state that represents the conformational changes due to sequential ligand binding [25]. Another study on the same receptor in Salipros showed the presence of a tetrameric assembly by cryo-EM and cryo-ET of secreted extracellular vesicles (Figure 4c) [15]. The authors suggest that this tetrameric state represents an assembly intermediate [15].

A series of structures of the human proton-activated chloride channel ASOR, determined at neutral or acidic pH in Salipros as well as the detergents glycodiosgenin (GDN) and Digitonin also highlight the

effect of a lipid environment on protein conformation. The channel can exist in resting, activated, and desensitized states. While the authors were able to capture the resting and activated states in detergent by altering the pH, the saposin-reconstituted sample contained both states at acidic pH, along with a previously reported desensitized state [16]. Another example is the  $\beta$ -barrel assembly machine (BAM complex) from *Escherichia coli*, where reconstitution of the complex into SapA revealed that the BamA subunit's beta barrel domain adopts a laterally closed conformation, in contrast to the structure of the detergent solubilized complex [26].

While reconstitution of an IMP in other nanodisc systems may have similar effects on protein structure and

function as described above, Salipros have the additional advantage of auto-tuning the nanodisc size to accommodate the target IMP [2]. This property not only simplifies the reconstitution procedure; it may also result in substantially improved reconstruction resolution in cases where transmembrane domains show substantial flexibility. An example of this is the structure of the mitochondrial calcium uniporter, which was determined to 3.8 Å resolution after reconstitution in Salipros, but only to 15 Å resolution when reconstituted in other nanodisc systems [17].

### Applications of Salipros beyond structural studies

Beyond their use in structure determination, Salipros have been proven to be valuable tools for studying the conformational changes and dynamics of membrane proteins using advanced techniques such as single-molecule Förster resonance energy transfer (smFRET) [27], nuclear magnetic resonance (NMR) [28], and mass spectrometry (MS) [29].

A key example is the dynamic study of the proton-coupled oligopeptide transporter, DtpA [27]. During a single transport cycle, this transporter transitions between at least three distinct conformational states. Using smFRET, researchers demonstrated that detergents tend to trap DtpA in a single conformational state, whereas in more native-like lipid environments, populations encompassing all functional states were observed. Furthermore, varying the lipid composition allowed for fine-tuning of the abundance of specific states [27].

In NMR studies, the stability and adaptability of Salipros have addressed long-standing challenges in membrane protein solubilization and spectral resolution. This size-tunable nanoparticle maintains a high lipid content to mimic realistic membrane environment while enabling stable, long-term solution NMR studies. This has been demonstrated with proteins like OmpX, pSR11, and  $\beta$ 1AR, the latter retaining functional interactions with binding partners [28]. The Salipro technology has also been used in mass spectrometry (MS) applications, in particular in the field of native MS. Zhou *et al.* demonstrated how Salipros maintain ferroportin's native structure and functionality during footprinting, achieving high sequence coverage [29].

### Further developments: direct extraction method

Conventionally, reconstitution into Salipros is achieved by mixing a detergent-solubilized membrane protein with SapA and lipids of choice. However, this approach requires the target protein to be purified and stable for an extended period under detergent-solubilized conditions. Prior information or optimization is necessary to

choose the ideal lipid composition to promote stabilization of an active membrane protein within the Salipro assembly. To overcome these obstacles, direct extraction of IMPs using Salipros, also known as DirectMX, was introduced to preserve native lipids bound to the IMP of choice [6]. However, the technique does not fully exclude the addition of detergents since a brief incubation step with digitonin or other mild detergents is required to disrupt the cellular membrane to allow for Salipro reconstitution (Figure 1b). However, after reconstitution, detergent is no longer necessary during subsequent steps. This reduces the time and efforts spent to identify a suitable detergent. DirectMX has been successfully applied in purifying the human solute carrier Excitatory Amino Acid Transporter 1 (EAAT1) [6] and in obtaining a high-resolution cryo-EM structure of the mouse ion channel Pannexin-1 (PANX1) [30]. For the latter, human and mouse PANX1 were directly isolated from Sf9 and HEK293 cells and purified to homogeneity. Moreover, the authors used His-tagged or biotinylated SapA to perform surface plasmon resonance measurements of ligand binding to reconstituted PANX1 [30]. This highlights the versatility of using Salipros beyond structure determination as a tool for biophysical analysis of IMPs where additional labeling could hamper activity measurements of the protein of interest.

### Conclusions and perspective

In recent years, the Salipro method has emerged as an attractive alternative to detergents and other nanodisc systems. In comparison to detergents, Salipros enable the structural and functional characterization of IMPs in a more lipid-like environment and preserve bound lipids. In addition, the recombinant production of Saposin A is simple and cost effective, and expression constructs are available. Salipros have been a great boon to structure determination by cryo-EM, as evidenced by the fact that all recently released Salipro structures were determined by cryo-EM (Supplementary Table S1) as opposed to X-ray crystallography or NMR [31–42]. They share these advantages over detergents with other nanodisc systems such as the Membrane Scaffold Proteins (MSPs). However, Salipros self-tune their size depending on the target, which simplifies sample preparation but might be detrimental for the conformational flexibility of the IMP in case the interaction of the IMP with the scaffold protein is too tight. With the advent of the DirectMX technology, saposin reconstitution can also bypass the need for detergent solubilization prior to classical Salipro reconstitution. They share this property with synthetic nanodisc reagents such as styrene maleic anhydride lipid particles (SMALPs) [43]. However, SMALPs and related reagents may not be suitable for every IMP target and may require expensive and time-consuming screening and optimization. Taken together, Salipros combine many of the desirable

properties of other IMP reconstitution methods due to their biochemical properties, versatility, and simplicity.

## Declaration of competing interest

The authors declare that they have no current competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Acknowledgements

We thank members of the Löw group for input and feedback on the manuscript. We thank the authors of the publications cited for their contributions to scientific discovery. We apologize to the community if we have overlooked any IMP structures determined in Salipros that were not cited.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.sbi.2025.103050>.

## Data availability

The PDB files used for this Mini-Review are stated. IMP structures reconstituted in Salipros with a PDB release date before December 2024 were considered.

## References

Papers of particular interest, published within the period of review, have been highlighted as:

- \* of special interest
- \*\* of outstanding interest

1. Günsel U, Hagn F: **Lipid nanodiscs for high-resolution NMR studies of membrane proteins.** *Chem Rev* 2022, **122**: 9395–9421, <https://doi.org/10.1021/acs.chemrev.1c00702>.
  2. Frauenfeld J, Löving R, Armache J-P, Sonnen AF-P, Guettou F, Moberg P, Zhu L, Jegerschöld C, Flayhan A, Briggs JAG, Garoff H, Löw C, Cheng Y, Nordlund P: **A saposin-lipoprotein nanoparticle system for membrane proteins.** *Nat Methods* 2016, **13**:345–351, <https://doi.org/10.1038/nmeth.3801>.
  3. Ahn VE, Faull KF, Whitelegge JP, Fluharty AL, Privé GG: **Crystal structure of saposin B reveals a dimeric shell for lipid binding.** *Proc Natl Acad Sci USA* 2003, **100**:38–43, <https://doi.org/10.1073/pnas.0136947100>.
  4. Popovic K, Holyoake J, Pomès R, Privé GG: **Structure of saposin A lipoprotein discs.** *Proc Natl Acad Sci USA* 2012, **109**: 2908–2912, <https://doi.org/10.1073/pnas.1115743109>.
  5. Kurgan KW, Chen B, Brown KA, Falco Cobra P, Ye X, Ge Y, Gellman SH: **Stable picodisc assemblies from saposin proteins and branched detergents.** *Biochemistry* 2021, **60**: 1108–1119, <https://doi.org/10.1021/acs.biochem.0c00924>.
  6. Lloris-Garcerá P, Klinter S, Chen L, Skynner MJ, Löving R, Frauenfeld J: **DirectMX – one-step reconstitution of membrane proteins from crude cell membranes into Salipro nanoparticles.** *Front Bioeng Biotechnol* 2020, **8**.
  7. Chojnacka W, Teng J, Kim JJ, Jensen AA, Hibbs RE: **Structural insights into GABAA receptor potentiation by quaalude.** *Nat Commun* 2024, **15**:5244, <https://doi.org/10.1038/s41467-024-49471-y>.
  8. Legesse DH, Fan C, Teng J, Zhuang Y, Howard RJ, Noviello CM, Lindahl E, Hibbs RE: **Structural insights into opposing actions of neurosteroids on GABAA receptors.** *Nat Commun* 2023, **14**: 5091, <https://doi.org/10.1038/s41467-023-40800-1>.
  9. Goswami U, Rahman MM, Teng J, Hibbs RE: **Structural interplay of anesthetics and paralytics on muscle nicotinic receptors.** *Nat Commun* 2023, **14**:3169, <https://doi.org/10.1038/s41467-023-38827-5>.
  10. Zhang K, Wu H, Hoppe N, Manglik A, Cheng Y: **Fusion protein strategies for cryo-EM study of G protein-coupled receptors.** *Nat Commun* 2022, **13**:4366, <https://doi.org/10.1038/s41467-022-32125-2>.
- This publication describes how saposin reconstitution of an important protein in the hedgehog signalling pathway revealed a previously unknown sterol binding site, as well as enabling the capture of a new conformation of the protein with potential consequences for its function within the pathway
11. Rahman MM, Basta T, Teng J, Lee M, Worrell BT, Stowell MHB, Hibbs RE: **Structural mechanism of muscle nicotinic receptor desensitization and block by curare.** *Nat Struct Mol Biol* 2022, **29**:386–394, <https://doi.org/10.1038/s41594-022-00737-3>.
  12. Flayhan A, Mertens HDT, Ural-Blimke Y, Martinez Molledo M, Svergun DI, Löw C: **Saposin lipid nanoparticles: a highly versatile and modular tool for membrane protein research.** *Struct. Lond. Engl.* 1993 2018, **26**:345–355.e5, <https://doi.org/10.1016/j.str.2018.01.007>.
  13. Guo J, Li S, Bai L, Zhao H, Shang W, Zhong Z, Maimaiti T, Gao X, Ji N, Chao Y, Li Z, Du D: **Structural transition of GP64 triggered by a pH-sensitive multi-histidine switch.** *Nat Commun* 2024, **15**:7668, <https://doi.org/10.1038/s41467-024-51799-4>.
  14. Nagamura R, Fukuda M, Kawamoto A, Matoba K, Dohmae N, Ishitani R, Takagi J, Nureki O: **Structural basis for oligomerization of the prokaryotic peptide transporter PeptSo2.** *Acta Crystallogr. Sect. F Struct. Biol. Commun.* 2019, **75**:348–358, <https://doi.org/10.1107/S2053230X19003546>.
  15. Introini B, Cui W, Chu X, Zhang Y, Alves AC, Eckhardt-Strelau L, Golusik S, Tol M, Vogel H, Yuan S, Kudryashev M: **Structure of tetrameric forms of the serotonin-gated 5-HT3A receptor ion channel.** *EMBO J* 2024, **43**:4451–4471, <https://doi.org/10.1038/s44318-024-00191-5>.
- By saposin reconstitution the authors were able to capture an assembly intermediate for this receptor, representing a previously unknown state which was confirmed by cryo-electron tomography studies of the receptor secreted in extracellular vesicles.
16. Wang C, Polovitskaya MM, Delgado BD, Jentsch TJ, Long SB: **Gating choreography and mechanism of the human proton-activated chloride channel ASOR.** *Sci Adv* 2022, **8**, eabm3942, <https://doi.org/10.1126/sciadv.abm3942>.
  17. Nguyen NX, Armache J-P, Lee C, Yang Y, Zeng W, Mootha VK, Cheng Y, Bai X-C, Jiang Y: **Cryo-EM structure of a fungal mitochondrial calcium uniporter.** *Nature* 2018, **559**:570–574, <https://doi.org/10.1038/s41586-018-0333-6>.
  18. Larsen ST, Dannersø JK, Nielsen CJF, Poulsen LR, Palmgren M, Nissen P: **Conserved N-terminal regulation of the ACA8 calcium pump with two calmodulin binding sites.** *J Mol Biol* 2024, **436**, 168747, <https://doi.org/10.1016/j.jmb.2024.168747>.
  19. Kim JJ, Gharpure A, Teng J, Zhuang Y, Howard RJ, Zhu S, Noviello CM, Walsh RM, Lindahl E, Hibbs RE: **Shared structural mechanisms of general anaesthetics and Benzodiazepines.** *Nature* 2020, **585**:303–308, <https://doi.org/10.1038/s41586-020-2654-5>.
  20. Zhu S, Sridhar A, Teng J, Howard RJ, Lindahl E, Hibbs RE: **Structural and dynamic mechanisms of GABAA receptor modulators with opposing activities.** *Nat Commun* 2022, **13**: 4582, <https://doi.org/10.1038/s41467-022-32212-4>.
  21. Fan C, Cowgill J, Howard RJ, Lindahl E: **Divergent mechanisms of steroid inhibition in the human P1 GABAA receptor.** *Nat Commun* 2024, **15**:7795, <https://doi.org/10.1038/s41467-024-51904-7>.
- In this publication, the authors identified a novel mechanism for subtype specific inhibition of the GABAA receptor by neurosteroid binding.
22. Hénault CM, Sun J, Therien JPD, daCosta CJB, Carswell CL, Labriola JM, Juranka PF, Baenziger JE: **The role of the M4 lipid-sensor in the folding, trafficking, and allosteric activation of**

- nicotinic acetylcholine receptors.** *Neuropharmacology* 2015, **96**:157–168, <https://doi.org/10.1016/j.neuropharm.2014.11.011>.
23. Rahman MM, Teng J, Worrell BT, Noviello CM, Lee M, Karlin A, Stowell MHB, Hibbs RE: **Structure of the native muscle-type nicotinic receptor and inhibition by Snake Venom Toxins.** *Neuron* 2020, **106**:952–962.e5, <https://doi.org/10.1016/j.neuron.2020.03.012>.
  24. Du D, Neuberger A, Orr MW, Newman CE, Hsu P-C, Samsudin F, Szweczak-Harris A, Ramos LM, Debela M, Khalid S, Storz G, Luisi BF: **Interactions of a Bacterial RND transporter with a transmembrane small protein in a lipid environment.** *Struct. Lond. Engl.* 1993 2020, **28**:625–634.e6, <https://doi.org/10.1016/j.str.2020.03.013>.
  25. Zhang Y, Dijkman PM, Zou R, Zandl-Lang M, Sanchez RM, Eckhardt-Strelau L, Köfeler H, Vogel H, Yuan S, Kudryashev M: **Asymmetric opening of the Homopentameric 5-HT<sub>3A</sub> serotonin receptor in lipid bilayers.** *Nat Commun* 2021, **12**:1074, <https://doi.org/10.1038/s41467-021-21016-7>.
  26. Xiao L, Han L, Li B, Zhang M, Zhou H, Luo Q, Zhang X, Huang Y: **Structures of the  $\beta$ -barrel assembly machine recognizing outer membrane protein substrates.** *FASEB J. Off. Publ. Fed. Am. Soc. Exp. Biol.* 2021, **35**, e21207, <https://doi.org/10.1096/fj.202001443RR>.
  27. Lasitzka-Male T, Bartels K, Jungwirth J, Wiggers F, Rosenblum G, Hofmann H, Löw C: **Membrane chemistry tunes the structure of a peptide transporter.** *Angew Chem Int Ed Engl* 2020, **59**: 19121–19128, <https://doi.org/10.1002/anie.202008226>.  
This publication highlights the important role of a lipid environment for an IMP. This transport protein has to undergo various conformational changes during its transport cycle. However detergent solubilization traps the protein in only one state, while Salipro reconstitution enabled capturing the protein in several conformations, and demonstrated that fine-tuning the lipid mixture could alter the relative abundance of each state.
  28. Chien C-TH, Helfinger LR, Bostock MJ, Solt A, Tan YL, Nietlispach D: **An adaptable phospholipid membrane mimetic system for solution NMR studies of membrane proteins.** *J Am Chem Soc* 2017, **139**:14829–14832, <https://doi.org/10.1021/jacs.7b06730>.
  29. Zhou F, Yang Y, Chemuru S, Cui W, Liu S, Gross M, Li W: **Footprinting mass spectrometry of membrane proteins: ferroportin reconstituted in saposin A picodiscs.** *Anal Chem* 2021, **93**:11370–11378, <https://doi.org/10.1021/acs.analchem.1c02325>.
  30. Drulyte I, Gutgsell AR, Lloris-Garcerá P, Liss M, Geschwindner S, Radjainia M, Frauenfeld J, Löving R: **Direct cell extraction of membrane proteins for structure–function analysis.** *Sci Rep* 2023, **13**:1420, <https://doi.org/10.1038/s41598-023-28455-w>.  
This publication describes the use of direct extraction using SapA for cryo-EM and SPR studies. Direct extraction from native membranes is a key benefit of saposins.
  31. Dai Y, Lee C-H: **Transport mechanism and structural pharmacology of human urate transporter URAT1.** *Cell Res* 2024, **34**:776–787, <https://doi.org/10.1038/s41422-024-01023-1>.
  32. Dalal V, Arcario MJ, Petroff JT, Tan BK, Dietzen NM, Rau MJ, Fitzpatrick JAJ, Brannigan G, Cheng WWL: **Lipid nanodisc scaffold and size alter the structure of a pentameric ligand-gated ion channel.** *Nat Commun* 2024, **15**:25, <https://doi.org/10.1038/s41467-023-44366-w>.
  33. Zhang J, Maksaev G, Yuan P: **Open structure and gating of the Arabidopsis mechanosensitive ion channel MSL10.** *Nat Commun* 2023, **14**:6284, <https://doi.org/10.1038/s41467-023-42117-5>.
  34. Chen H, Ahmed S, Zhao H, Elghobashi-Meinhardt N, Dai Y, Kim JH, McDonald JG, Li X, Lee C-H: **Structural and functional insights into Spns2-mediated transport of Sphingosine-1-Phosphate.** *Cell* 2023, **186**:2644–2655.e16, <https://doi.org/10.1016/j.cell.2023.04.028>.
  35. Cowgill J, Fan C, Haloi N, Tobiasson V, Zhuang Y, Howard RJ, Lindahl E: **Structure and dynamics of Differential ligand binding in the human  $\rho$ -Type GABAA receptor.** *Neuron* 2023, **111**:3450–3464.e5, <https://doi.org/10.1016/j.neuron.2023.08.006>.
  36. Xu B, Zhang M, Zhang B, Chi W, Ma X, Zhang W, Dong M, Sheng L, Zhang Y, Jiao W, Shan Y, Chang W, Wang P, Wen S, Pei D, Chen L, Zhang X, Yan H, Ye S: **Embigin facilitates monocarboxylate transporter 1 localization to the plasma membrane and transition to a decoupling state.** *Cell Rep* 2022, **40**, 111343, <https://doi.org/10.1016/j.celrep.2022.111343>.
  37. Noviello CM, Kreye J, Teng J, Prüss H, Hibbs RE: **Structural mechanisms of GABAA receptor autoimmune encephalitis.** *Cell* 2022, **185**:2469–2477.e13, <https://doi.org/10.1016/j.cell.2022.06.025>.
  38. Kehlenbeck DM, Traore DAK, Josts I, Sander S, Moulin M, Haertlein M, Prevost S, Forsyth VT, Tidow H: **Cryo-EM Structure of MsbA in saposin-lipid nanoparticles (Salipro) provides insights into nucleotide coordination.** *FEBS J* 2022, **289**: 2959–2970, <https://doi.org/10.1111/febs.16327>.
  39. Ornik-Cha A, Wilhelm J, Kobyłka J, Sjuets H, Vargiu AV, Mallocci G, Reitz J, Seybert A, Frangakis AS, Pos KM: **Structural and functional analysis of the promiscuous AcrB and AdeB efflux pumps suggests different drug binding mechanisms.** *Nat Commun* 2021, **12**:6919, <https://doi.org/10.1038/s41467-021-27146-2>.
  40. Noviello CM, Gharpure A, Mukhtasimova N, Cabuco R, Baxter L, Borek D, Sine SM, Hibbs RE: **Structure and gating mechanism of the A7 nicotinic acetylcholine receptor.** *Cell* 2021, **184**: 2121–2134.e13, <https://doi.org/10.1016/j.cell.2021.02.049>.
  41. Gharpure A, Teng J, Zhuang Y, Noviello CM, Walsh RM, Cabuco R, Howard RJ, Zaveri NT, Lindahl E, Hibbs RE: **Agonist selectivity and ion permeation in the A3 $\beta$ 4 ganglionic nicotinic receptor.** *Neuron* 2019, **104**:501–511.e6, <https://doi.org/10.1016/j.neuron.2019.07.030>.
  42. Kintzer AF, Green EM, Dominik PK, Bridges M, Armache J-P, Deneka D, Kim SS, Hubbell W, Kossiakoff AA, Cheng Y, Stroud RM: **Structural basis for activation of voltage sensor domains in an ion channel TPC1.** *Proc Natl Acad Sci* 2018, **115**, <https://doi.org/10.1073/pnas.1805651115>.
  43. Knowles TJ, Finka R, Smith C, Lin Y-P, Dafforn T, Overduin M: **Membrane proteins solubilized intact in lipid containing nanoparticles bounded by styrene maleic Acid copolymer.** *J Am Chem Soc* 2009, **131**:7484–7485, <https://doi.org/10.1021/ja810046q>.